arXiv:hep-lat/9712023v1 19 Dec 1997

Submitted to Phys. Rev. Lett.

Temperature dependence of instantons in QCD

M.-C. Chu(1), S. M. Ouellette(2), S. Schramm(3), and R. Seki(4,5)
(1) *Department of Physics, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong*
(2) *Lauritsen Laboratory, Caltech 452-48, Pasadena, CA 91125, U.S.A.*
(3) *GSI, Darmstadt, Germany*
(4) *W. K. Kellogg Radiation Laboratory, Caltech 106-38, Pasadena, CA 91125, U.S.A.*
(5) *Department of Physics, California State University, Northridge, CA 91330, U.S.A.*

(February 1, 2008)

Abstract

We investigate the temperature dependence of the instanton contents of gluon fields, using unquenched lattice QCD and the cooling method. The instanton size parameter deduced from the correlation function decreases from 0.44fm below the phase-transition temperature T_c (≈ 150MeV) to 0.33fm at 1.3 T_c. The instanton charge distribution is Poissonian above T_c, but it deviates from the convoluted Poisson at low temperature. The topological susceptibility decreases rapidly below T_c, showing the apparent restoration of the $U(1)_A$ symmetry already at $T \approx T_c$.

Typeset using REVTEX

Among the four global symmetries associated with quantum chromodynamics (QCD), two chiral symmetries, $SU(N_f)_A$ and $U(1)_A$, must be broken as the observed hadron spectrum suggests. At high temperature, it is expected that the two symmetries are restored. The question as to how they are restored is an important issue, especially in relation to the QCD deconfinement phase transition (or maybe a crossover), and is recently attracting much attention [1–4]. The deconfinement phase transition is thought to have occurred in the evolution of the early universe and is expected to generate a new phase of matter, the quark-gluon plasma (QGP), which is under investigation in relativistic heavy-ion collisions.

Instantons, tunneling between topologically different gauge vacua, are believed to play crucial roles in the breaking and restoration of the chiral symmetries, and perhaps also in the deconfinement phase transition [2]: The $SU(N_f)_A$ symmetry manifests itself in the Goldstone mode through chiral condensates generated dynamically by instantons. A similar mechanism may be at work in the formation of hadronic bound states, and the $SU(N_f)_A$ restoration and the deconfinement are thus closely related to each other. The $U(1)_A$ symmetry is spontaneously broken by the chiral anomaly driven by instantons. At high temperatures, the restoration of the symmetries takes place because instanton amplitudes are suppressed through Debye screening, which induces the gluons to be (electrically) massive.

How realistic is this instanton mechanism, and what is the precise dynamics of the instanton participation? The best means to answer the questions is the lattice gauge theory [5], which provides a systematic, nonperturbative framework to study the equilibrium properties of QCD at both zero and finite temperatures. Whereas direct comparison between lattice results and experiments is not always possible, lattice QCD often motivates and constrains phenomenological models, which connect the QCD with experiments. Here, the use of *unquenched* QCD must be essential, since the instanton mechanism is expected to be affected by the presence of the dynamical quarks.

In this letter, we report unquenched lattice QCD results on the temperature dependence of the instanton contents of gluons. For the first time, we find curious temperature dependence suggesting that instantons are seemingly correlated below the $SU(N_f)_A$ phase transition, $T_c \approx 150$MeV. The nature and the details of the correlation are yet unclear, however, and so is the question of whether it corresponds to any of the dynamical models [2]. We also find that the $U(1)_A$ symmetry appears to be largely already restored at T_c. Note that because of the cooling method we used, we observed neither small instantons nor instanton molecules.

The calculation is carried out for $16^3 \times N_t$ lattices with $N_t = 4, 6, 8, 10, 12, 14$, and 16 for $m_q a = 0.0125$. The temperature is varied by changing N_t, with $\beta = 5.54$ fixed. The susceptibility and the topological-charge correlation are observed to be stable with respect to the number of cooling steps. We find that the topological susceptibility decreases slowly at low temperature as the temperature increases, but a rapid suppression sets in around $T = T_c$, so that, compared to the value at $T = 0$, it is already down by an order of magnitude at T_c and becomes only about 1% at $1.3T_c$. Accompanying this rapid change in χ_t is a transition of the instanton size from $\rho \approx 0.44$fm at low temperature to 0.33fm at $1.3T_c$. The distribution of instanton charges is Poissonian above T_c, but it deviates from the convoluted Poisson at low temperatures.

The instanton content of the gauge fields is monitored by the the topological-charge density, which can be measured on the lattice with

$$Q(x_n) = -\frac{1}{32\pi^2}\epsilon_{\alpha\beta\gamma\delta}\text{Re Tr}\left[U_{\alpha\beta}(x_n)U_{\gamma\delta}(x_n)\right] \ , \tag{1}$$

where $U_{\alpha\beta}$ is the product of the link variables around a plaquette in the $\alpha-\beta$ plane. A direct calculation of Q from gauge configurations is plagued with poor statistics and with lattice artifacts associated with discretization. Teper *et al.* [6] have shown that the topological charge can be reliably extracted using the cooling method, in which one applies a certain number of smoothing steps to each configuration in order to minimize the action locally. The ultraviolet fluctuations are quickly suppressed by these cooling steps, and only the relatively stable instantons are left in the configuration, making it easy to extract the topological charge. Details of the cooling method are discussed in [6,7].

The topological susceptibility is then given as the fluctuations of the topological charge:

$$\chi_t \equiv \frac{1}{N_t N_s^3 a^4}\langle\left(\sum_n Q(x_n)\right)^2\rangle \ , \tag{2}$$

where $\langle ... \rangle$ indicates configuration averaging, N_t and N_s are the number of sites in the temporal and spatial direction, respectively, and a is the lattice spacing.

In order to examine the details of the topological-charge distribution, we also calculate the topological charge-density correlation function [7],

$$C_Q(x) = \langle\sum_y Q(y)Q(x+y)\rangle/\langle\sum_y Q^2(y)\rangle \ . \tag{3}$$

One can compare this correlation function to a convolution of an isolated instanton topological-charge density

$$Q_\rho(x) = \frac{6}{\pi^2\rho^4}\left(\frac{\rho^2}{x^2+\rho^2}\right)^4 \ . \tag{4}$$

Zero-temperature calculations using the cooling method [7] show that after about 50 cooling steps, the gauge fields are dominated by large, isolated instantons, whose profiles agree well with Eq. (4).

To generate the unquenched gauge configurations, we adopted and modified a code written by the MILC collaboration [9], using its option of dynamical Kogut-Susskind fermions. Forty configurations are generated for each of $N_t = 12, \ 14, \ 16$, and 100 configurations for each of $N_t = 4, \ 6, \ 8, \ 10$. In each case, a molecular-dynamics time step of $dt = 0.02$ is used, and the first 300 time units (43 for $N_t = 12, \ 14, \ 16$) from a hot start are used as thermalization. The configurations were separated by 3 time units for $N_t = 12, \ 14, \ 16$; 5 time units for $N_t = 4, \ 6, \ 10$; and 10 time units for $N_t = 8$.

These configurations are then cooled, using the standard procedure as used in [7], for 200 steps, saving every 20 steps. We have checked to see that both the topological susceptibility and the topological-charge correlation function settle to stable values by 100 cooling steps. The results reported here are based on those extracted after 120 cooling steps.

The temperature of a gauge configuration is given by $T = 1/(N_t a)$ and can be varied by changing either N_t, or a (= the lattice spacing). a is a function of the lattice inverse coupling β, and can be calculated by the perturbation theory, but only for a narrow range of β (and hence to T) in the asymptotic scaling regime. In order to obtain the temperature dependence reliably, we choose to vary the temperature by varying N_t, keeping a and β fixed. In this way, the uncertainty in a enters only as an overall constant factor, which little affects the functional relation of the χ_t and T. The price we pay is that the calculation can be done only for a limited number of temperatures corresponding to the discrete values of N_t. This is especially problematic at high temperatures. The phase transition with the parameters we use occurs at $N_t = 8$ and a temperature of $T_c = 150(9)$ MeV [9], giving $a \approx 0.17$fm.

Figure 1 shows the topological susceptibility as a function of the temperature. The solid curve in the figure is the PCAC suppression that is due to soft pion gas [10], which is believed to be relevant for low temperature $T << T_c$:

$$\chi_t(T) = \chi_t(T=0)\left(1 + cT^2/{F_\pi}^2\right) \ . \tag{5}$$

Equation 5 fits our data very well up to $T = T_c$, but with $c = -0.36$, a value that is well outside the allowed range based on the PCAC argument $-1/6 \leq c \leq 1/6$ [10]. At higher temperatures, the suppression of the susceptibility follows a pattern different from the PCAC prediction.

Also shown in Fig. 1 is the perturbative Debye-screening prediction (for $N_c = 3,\ N_f = 2$) [11]:

$$\chi_t(T) = \chi_t(T=0)\left(1 + \lambda^2/3\right)^{\frac{7}{6}} \exp\left[-8\lambda^2/3 - 14\alpha\left(1 + \gamma\lambda^{-\frac{3}{2}}\right)^{-8}\right] \ , \tag{6}$$

where $\lambda \equiv \pi\rho T$, $\alpha = 0.01289764$, and $\gamma = 0.15858$. The dot-dashed and dashed curves correspond to $\rho = 0.44$ and 0.33 fm, respectively. Equation 6 is not expected to be valid below T_c, but should become relevant at high temperatures. We observe that the curve for $\rho = 0.44$ fm fits the four lowest temperature points quite well, giving an alternative suppression model to PCAC at low temperatures, and that the curve for $\rho = 0.33$ fm fits the high temperature points quite well. This observation suggests that the Debye-screening mechanism may be already operative for $T < T_c$, though the dominant instanton size parameter changes suddenly across T_c. Unfortunately, our $N_t = 4$ ($T = 2T_c$) result is rather uncertain because of severe finite-size effects and because not one single instanton is found in our 100 configurations. Also, our method of temperature variation by varying N_t provides only sparse data in the high-temperature regime. The temperature dependence of the Debye-screening predictions must be explicitly verified by further calculations.

The general features of the temperature dependence of the χ_t are the same in the quenched calculation [8] and in the present unquenched calculation, but detailed features differ significantly. The unquenched χ_t decreases more slowly at low temperature, but goes down much more rapidly around T_c. The χ_t is suppressed by almost 90 % at $T = T_c$ in the unquenched case, while it is only 50 % in the quenched case. That is, the suppression of the instanton amplitudes is enhanced by the dynamical quarks. The rapid decrease in χ_t continues above T_c, so that only about 1% remains at $1.3T_c$ in the unquenched, and about 10% in the quenched calculation. This temperature dependence of χ_t suggests that the window of temperatures above T_c in which $U_A(1)$ is broken is *narrowed* by the presence of dynamical quarks, and that $U_A(1)$ is practically restored at about T_c. One must take caution, however, in applying lattice results on the $U(1)_A$ symmetry, because the symmetry is explicitly broken in the conventional lattice and the process of approaching the continuum limit involves uncontrolled subtlety [3]. On the basis of the present calculation with $m_q a = 0.0125$, we draw no conclusion on the relationship between T_c and the $U(1)_A$ symmetry breaking temperature, and thus on whether the phase transition is of the first or second order [12,13].

Figure 2 shows the temperature dependence of the topological charge correlation functions in comparison to the profile of an isolated instanton, Eq.(4). In the quenched calculations, $C_Q(r)$ is practically independent of the temperature, following remarkably closely to the profile, though hints of a transition to a smaller size parameter have been observed above T_c [8]. The present unquenched calculation confirms the same trend below T_c, but it clearly shows a rapid transition to a smaller size around T_c. For clarification, we show in Fig. 2 the temperature dependence only around T_c: the size parameter, ρ, is approximately 0.44 fm at $0.75T_c$, shrinks rapidly around T_c, and becomes about 0.33fm at $1.3T_c$. It is interesting to note that the low-temperature value of ρ agrees fairly well with that used in the instanton liquid model, while the high-temperature value agrees with the dilute-gas model and also with the quenched value [2].

The distribution of instanton charges in our configuration samples is shown as histograms for each N_t in Fig. 3. Whereas in the quenched case, these distributions agree with the convolued Poisson distribution [8], the unquenched results show significant deviations from this Poissonian, with curious suppression of the distribution around the zero topological charge ($Q = 0$) for all N_t below the phase transition. The dynamical quarks have apparently induced non-trivial correlations among the instantons at low temperatures. Above the phase transition, the distributions become Poissonian, suggesting a return to the dilute-gas ensemble.

The instantons we have discussed so far in this letter are all of the topological charge of the unit magnitude. We note that we have observed occassionally the instantons of the topological charge of the magnitude two, which are rarely treated in any dynamical model.

In conclusion, our investigation of the temperature dependence of the instanton contents of unquenched gluon fields yields the following results: The topological susceptibility decreases rapidly around the phase-transition temperature $T_c \approx 150$MeV; at $T = T_c$, it is suppressed by one order of magnitude compared to its zero-temperature value, and at $T = 1.3T_c$, it is reduced by another order of magnitude. The temperature dependence of the topological susceptibility is consistent with a PCAC prediction for temperatures up to T_c, though with a large coefficient c. It also agrees with a Debye-screening model, but with a sudden shrinking of the instanton size parameter at around T_c. The topological charge correlation function also changes rapidly around T_c, which can be characterized by a transition of the dominant instanton size parameter from 0.44fm at low temperatures to 0.33fm at 1.3 T_c. The distribution of the instanton charges suggests the existence of nontrivial correlations among the instantons at low temperatures, which disappear above T_c. Our results indicate that the $U_A(1)$ symmetry is apparently restored largely at $T \approx T_c$.

This work is in part based on the MILC collaboration's public lattice gauge theory code. We thank C. DeTar and C. McNeile for their technical assistance. We acknowledge San Diego Supercomputer Center for providing Intel Paragon and Cray T3E computer resources. This research is partially supported by the U. S. National Science Foundation under grants PHY88-17296 and PHY90-13248 at Caltech, the U.S. Department of Energy under grant DE-FG03-87ER40347 at CSUN, and the Institute of Mathematical Sciences at the Chinese

University of Hong Kong. M. C. acknowledges the support of a summer research grant and a direct grant (2060105) of the Chinese University of Hong Kong.

REFERENCES

[1] E. Shuryak, Comments Nucl. Part. Phys. **21**, 235 (1994).

[2] T. Schäfer and E. V. Shuryak, University of Washington preprint, hep-ph/9610451, 1997; to appear in Rev. Mod. Phys.

[3] C. Bernard *et al.*, Phys. Rev. Lett. **78**, 598 (1997).

[4] G. Boyd, F. Karsch, E. Laermann, and M. Oevers, IFUP-TH 40/96 and BI-TP 96/27 (hep-lat/9607046) 1996; P. deForcrand, M. G. Pérez, and I-O Stamatescu, hep-lat/9701012, 1997: and P. deForcrand, M. G. Pérez, J. E. Hetrick, and I-O Stamatescu, hep-lat/9710001, 1997.

[5] I. Montvay and G. Münster, *Quantum Fields on Lattice* (Cambridge Univ. Press, Cambridge, 1994), and references therein.

[6] M. Teper, Phys. Lett. **B171**, 81 (1986); and J. Hoek, M. Teper, and J. Waterhouse, Nucl. Phys. **B288**, 589 (1987).

[7] M.-C. Chu, J. M. Grandy, S. Huang, and J. W. Negele, Phys. Rev. **D49**, 6039 (1994); Phys. Rev. Lett. **70**, 225 (1993); M.-C. Chu and S. Huang, Phys. Rev. **D45**, 2446 (1992).

[8] M. -C. Chu and S. W. Schramm, Phys. Rev. **D51**, 4580 (1995).

[9] S. Gottlieb *et al.*, Phys. Rev. **D47**, 3619 (1993).
See http://physics.indiana.edu/s̃g/milc.html.

[10] E. Shuryak and M. Velkovsky, Phys. Rev. **D50**, 3323 (1994).

[11] R. D. Pisarski and L. G. Yaffe, Phys. Lett. **97B**, 110 (1980).

[12] R. D. Pisarski and F. Wilczek, Phys. Rev. **D29**, 338 (1984).

[13] F. Wilczek, Int. J. Mod. **D3** Suppl., 63 (1994); and K. Rajogopal and F. Wilczek, Nucl. Phys. **B399**, 395 (1993).

FIGURES

FIG. 1. Topological susceptibility as a function of temperature. The data points correspond to our unquenched-QCD-plus-cooling results with $N_t = 6, 8, 10, 12, 14, 16,$ and $a = 0.17$fm is assumed. The transition temperature T_c is approximately 150 MeV for our calculations with $\beta = 5.54, ma = 0.0125, N_x = 16$. The dashed curve is the prediction of a PCAC model, Eq. (5), with $c = -0.36$; and the solid and dot-dashed curves of a Debye-screening model, Eq. (6), with the instanton size parameter, $\rho = 0.44$ and 0.33 fm, respectively.

FIG. 2. Topological charge correlation functions across the phase-transion temperature ($T_c = 150$MeV) from the same calculations as in Fig. 1.

FIG. 3. Histograms showing the distribution of topological charge in the configurations for $N_t = 6, 8, 10, 12, 14$ and 16. Q is the magnitude of the number of topological charges in a configuration, and $N(Q)$ is the frequency. The dots in the figure are the convoluted Poisson distributions fitted to the histograms.



SV: c=−0.36
PY: ρ=0.44 fm
PY: ρ=0.33 fm
10^−4
10^−5
10^−6
0.0
100.0
200.0
T (MeV)
X_t(T)a^4



△ T = 0.75 T_c
× T = T_c
□ T = 1.3 T_c
ρ = 0.44 fm
ρ = 0.33 fm
$C_Q(r)$
r/a
1.0
0.6
0.2
−0.2
0.0
2.0
4.0
6.0


N(Q)
N(Q)
N(Q)
Q
Q
N_t=6
N_t=8
N_t=10
N_t=12
N_t=14
N_t=16
0
50
100
150
0
1
0
20
40
60
80
0
1
2
3
0
10
20
30
40
0
2
4
6
8
0
5
10
15
0
2
4
6
8
10
0
5
10
15
0
5
10
15
0
2
4
6
8
10
0
5
10
15